

Reduce one of your biggest costs while achieving superior results



Typical Share of Landscape Budget*

Minimal Investment

- For a small premium per square foot, you can significantly reduce mowing time and cost

Maximum Value

- Mowing cost savings estimated at \$1,500 per acre, per year—every year*

*Landscape budget percentages based on Scotts' proprietary market research. Based on estimated \$60-\$200 cost per acre to mow.



For more information, visit:
ProVistaTurf.com

©2018 The Scotts Company LLC. World rights reserved. 19PR02



Have a Premium Landscape with Significantly Less Mowing and Cost

Available from
JASPERSON SOD
Spring 2021




Scotts ProVista Use and Care Instructions



Cost Effective

- Scotts® ProVista™ grows slower than conventional Kentucky Bluegrass, saving valuable mowing time and reducing maintenance costs



Complete Weed Control

- Easily combat weeds, without damaging turf
- Beautiful green, uniform, dense appearance

Sustainable

- Reduced mowing means less carbon emissions
- Reduced Inputs: stays green with less fertilizer and herbicide

Half the Mowing

- Save valuable time and money
- No change in mowing heights required

Follow Standard Cool Season Turf Maintenance Program

- Refer to your state extension site for best management practices
- No special care needed vs. conventional Kentucky Bluegrass

An Added Tool in Your Toolbox - for Invasive, Grassy Weeds

How to use Glyphosate on Scotts ProVista

- Only to be used on Scotts ProVista turf; will kill conventional grass.
- Only apply if alternative active ingredients were not effective or are not available for existing weed pressure.
- Utilize pre-emergent herbicides for weed prevention to minimize the need for broadcast glyphosate applications.
- Scotts ProVista turfgrass that is healthy and actively growing under normal growing conditions is tolerant to glyphosate. Injury may occur when applied under stresses, such as, but not limited to heat, drought, shade, salinity, disease, insect pressure, or cold (below 50°F within 10 days of an application).

- For establishment, a single broadcast glyphosate application should occur no sooner than one month after sodding and be tank mixed with a pre-emergence herbicide that is safe during the establishment period.
- When spraying glyphosate, tank mix other post-emergent herbicides with efficacy on the target weed to ensure multiple modes of action are applied.
- Apply at 32 fl oz per acre (or 0.75 fl oz per 1,000 ft²) in carrier volume of 40 gal per acre (or 1 gal per 1,000 ft²) with a maximum of 3 broadcast applications per calendar year.
- For difficult to control weeds, apply 48 fl oz per acre (or 1.125 fl oz per 1,000 ft²) in carrier volume of 40 gal per acre (or 1 gal per 1,000 ft²) with a maximum of 2 broadcast applications per calendar year.

